

# Maps of Surface Groups to Finite Groups with No Simple Loops in the Kernel

by

Charles Livingston

**Abstract:** Let  $F_g$  denote the closed orientable surface of genus  $g$ . What is the least order finite group,  $G_g$ , for which there is a homomorphism  $\psi: \pi_1(F_g) \rightarrow G_g$  so that no nontrivial simple closed curve on  $F_g$  represents an element in  $\text{Ker}(\psi)$ ? For the torus it is easily seen that  $G_1 = Z_2 \times Z_2$  suffices. We prove here that  $G_2$  is a group of order 32 and that an upper bound for the order of  $G_g$  is given by  $g^{2g+1}$ . The previously known upper bound was greater than  $2^{g^{2g}}$ .

For any compact surface  $F$  there exists a finite group  $G$  and a homomorphism  $\psi: \pi_1(F) \rightarrow G$  such that no nontrivial element in the kernel of  $\psi$  can be represented by a simple closed curve. Such a homomorphism is said to have *nongeometric kernel*. Casson, Gabai, and Skora [5] have each constructed examples of this (see Section 2 for details). The presence of such examples raises a variety of questions relating to the characterization of the finite groups that can occur in this way. This paper addresses the problem of determining the relationship between the genus of  $F$  and the order of  $G$ . In the case that  $F$  is a torus a complete analysis is straightforward. For instance, the natural projection  $\psi: \pi_1(F) \rightarrow H_1(F; Z_2) \cong Z_2 \times Z_2$  has nongeometric kernel.

Our first result concerns the genus 2 closed orientable surface,  $F_2$ . Casson's construction yields a group of order  $2^{38}$ . Skora reduced this order considerably by producing a group of order  $2^9$ . In Section 3 a group of order  $2^5 = 32$ ,  $G_2$ , is constructed for which there is a homomorphism  $\psi_2: \pi_1(F_2) \rightarrow G_2$  having nongeometric kernel. In Section 4 it is proved that no such example can be constructed using a group of order less than 32.

The example in Section 3 is generalized to construct examples for arbitrary genus surfaces in Section 5. The order of the groups constructed is quite small compared to previously constructed examples. As the examples directly generalize the minimal genus 2 example, there is the possibility that they are minimal as well.

*Acknowledgements* Thanks are due to Allan Edmonds for pointing out the proof of Theorem 4.2. The work in Section 5 was motivated by discussions with Dennis Johnson.

## 1 Notation and Conventions

Throughout this paper all surfaces will be closed and orientable. References to basepoints for the fundamental group of a space are omitted. Since the property of being in the kernel of a homomorphism depends only on the conjugacy class of an element, such omissions will not affect the arguments.

By a *simple loop* on a surface we mean an embedding of the circle  $S^1$ .

We will say that a homomorphism  $\psi: \pi_1(F) \rightarrow G$  has *geometric kernel* if some nontrivial element in the kernel can be represented by a simple loop. Otherwise  $\psi$  has *nongeometric kernel*.

## 2 Basic Examples

In this section a procedure of Casson is used to construct for each surface  $F$  a finite group  $G$  and a surjective homomorphism  $\psi: \pi_1(F) \rightarrow G$  such that  $\psi$  has nongeometric kernel. The orders of the groups involved is computed for contrast with the examples produced in Section 5.

The statement that  $\psi$  has nongeometric kernel can be reinterpreted in terms of covering spaces as

follows. Corresponding to  $\text{Ker}(\psi)$  there is a connected regular covering space  $p: \tilde{F} \rightarrow F$  with  $p_*(\pi_1(\tilde{F})) = \text{Ker}(\psi)$ . An element in  $\pi_1(F)$  is in  $\text{Ker}(\psi)$  if and only if when represented by a closed path, the path lifts to a closed path in  $\tilde{F}$ . Hence a simple loop on  $F$  represents an element in  $\text{Ker}(\psi)$  if and only if it can be lifted to a simple loop in  $\tilde{F}$ . Conversely, if  $p: \tilde{F} \rightarrow F$  is a regular covering space with the property that no nontrivial simple loop on  $F$  lifts to  $\tilde{F}$  then the natural projection  $\psi: \pi_1(F) \rightarrow \pi_1(F)/p_*(\pi_1(\tilde{F}))$  has nongeometric kernel.

**Construction** Given a surface  $F$ , construct the covering space  $p: \tilde{F} \rightarrow F$  corresponding to the kernel of the projection  $\pi_1(F) \rightarrow H_1(F; Z_2)$ . Simple nonseparating loops on  $F$  represent generators of  $H_1(F; Z_2)$  and hence do not lift to  $\tilde{F}$ . Nontrivial separating simple loops do lift, but each preimage on  $\tilde{F}$  is nonseparating on  $\tilde{F}$ .

Now construct the covering  $q: \bar{F} \rightarrow \tilde{F}$  corresponding to the kernel of the projection  $\pi_1(\tilde{F}) \rightarrow H_1(\tilde{F}; Z_2)$ . As no nonseparating simple loop on  $\tilde{F}$  lifts to  $\bar{F}$  it is apparent that no nontrivial simple loop on  $F$  lifts to  $\bar{F}$ .

It remains to show that the covering  $p \circ q: \bar{F} \rightarrow F$  is regular; that is, that  $p_* \circ q_*(\pi_1(\bar{F}))$  is normal in  $\pi_1(F)$ . Observe that  $q_*(\pi_1(\bar{F}))$  is a characteristic subgroup of  $\pi_1(\tilde{F})$  and  $p_*(\pi_1(\tilde{F}))$  is a characteristic subgroup of  $\pi_1(F)$ . Since a characteristic subgroup of a characteristic subgroup is characteristic,  $p_* \circ q_*(\pi_1(\bar{F}))$  is characteristic in  $\pi_1(F)$ , and is hence normal.

**Order of  $\pi_1(F)/\langle p_* \circ q_*(\pi_1(\bar{F})) \rangle$**

The order of this finite group is equal to the degree of the covering  $p \circ q$ . Suppose that  $F$  is of genus  $g$ . The Euler characteristic of  $F$  is  $2 - 2g$ . Since  $\tilde{F}$  is a  $2^{2g}$  fold cover of  $F$ , the Euler characteristic of  $\tilde{F}$  is  $2^{2g}(2 - 2g)$ . The genus of  $\tilde{F}$  is  $\frac{1}{2}(2 - 2^{2g}(2 - 2g)) = \frac{1}{2}((g - 1)2^{2g+1} + 2) = g'$ . The covering  $q: \bar{F} \rightarrow \tilde{F}$  is of degree  $2^{2g'}$ . The degree of  $p \circ q: \bar{F} \rightarrow F$  is the product of these two degrees:  $2^{2g'}2g = 2^{(g-1)2^{2g+1}+2+2g}$ .

**Note** The construction of Gabai differs considerably from the one above. He notes that every simple curve is in the complement of some index three subgroup of  $\pi_1(F)$ ; nonseparating curves are not in the kernel of some map to  $Z_3$  and separating curves are mapped to a 3-cycle in the third symmetric group,  $S_3$ , under some homomorphism and hence map to the complement of an index three subgroup of  $S_3$ . Hence the quotient map  $\pi_1(F) \rightarrow \pi_1(F)/H$  has nongeometric kernel, where  $H$  is the intersection of all index three subgroups of  $\pi_1(F)$ . We have been unable to find reasonable bounds on the size of this quotient.

### 3 A small genus 2 example

The groups constructed in the previous section are of very large order. If  $F$  is of genus 2, the corresponding group is of order  $2^{38}$ . This section presents a description of a group of order 32,  $G_2$ , and a homomorphism  $\psi_2$  of the fundamental group of the genus 2 surface to  $G_2$ , such that  $\psi_2$  has a nongeometric kernel. The next section contains a proof that this example is minimal.

For the remainder of this section  $F$  will denote a genus 2 surface.

**Construction of  $G_2$**  For our purposes, the easiest way to describe  $G_2$  is as follows. Define a group structure on the set  $(Z_2)^4 \times Z_2$  by defining the product by

$$(a_1, b_1, a_2, b_2, \epsilon)(a'_1, b'_1, a'_2, b'_2, \epsilon') = (a_1 + a'_1, b_1 + b'_1, a_2 + a'_2, b_2 + b'_2, \epsilon + \epsilon' + b_1a'_1 + b_2a'_2).$$

The operations within the parenthesis are addition and multiplication in  $Z_2$ . The verification that this defines a group structure can be done by a direct calculation, which is left to the reader. The group is denoted  $32_{42}$  in [6], and  $\Gamma_5 a_1$  in the notation of [3].

An essential calculation for later purposes is that of commutators in  $G_2$ . A direct computation yields

$$[(a_1, b_1, a_2, b_2, \epsilon), (a'_1, b'_1, a'_2, b'_2, \epsilon')] = (0, 0, 0, 0, (b_1a'_1 - b'_1a_1) + (b_2a'_2 - b'_2a_2)). \quad (1)$$

From this it is apparent that both the center and commutator subgroup of  $G_2$  consists of the set  $(0, 0, 0, 0) \times Z_2$ . The abelianization of  $G_2$  is  $(Z_2)^4$ , given by the projection  $(Z_2)^4 \times Z_2 \rightarrow (Z_2)^4 \times \{0\}$ .

**Construction of  $\psi_2$**  Let  $\{x_1, y_1, x_2, y_2\}$  be a standard generating set of  $\pi_1(F)$  so that  $\pi_1(F)$  has presentation  $\langle x_1, y_1, x_2, y_2, [x_1, y_1][x_2, y_2] = 1 \rangle$ . This set projects to a standard symplectic basis of  $H_1(F; Z_2)$ ,  $\{|x_1|, |y_1|, |x_2|, |y_2|\}$

Define  $\psi_2: \pi_1(F) \rightarrow G_2$  be setting:

$$\psi_2(x_1) = (1, 0, 0, 0) \times (0),$$

$$\psi_2(y_1) = (0, 1, 0, 0) \times (0),$$

$$\psi_2(x_2) = (0, 0, 1, 0) \times (0),$$

$$\psi_2(y_2) = (0, 0, 0, 1) \times (0).$$

Using (1) it is easily verified that this gives a well defined surjective representation.

The key observation is that  $\psi_2$  has the following property: if  $\omega_1$  and  $\omega_2$  are elements of  $\pi_1(F)$ , then

$$[\psi_2(\omega_1), \psi_2(\omega_2)] = (0, 0, 0, 0) \times (|\omega_1| \cap |\omega_2|), \quad (2)$$

where  $|\omega_1| \cap |\omega_2|$  is the  $Z_2$  intersection number of the classes in  $H_1(F; Z_2)$ , represented by  $\omega_1$  and  $\omega_2$ . This follows from (1) along with the fact that if  $(a_1, b_1, a_2, b_2)$  and  $(a'_1, b'_1, a'_2, b'_2)$  are classes in  $H_1(F; Z_2)$ , then  $(a_1, b_1, a_2, b_2) \cap (a'_1, b'_1, a'_2, b'_2) = (b_1 a'_1 - b'_1 a_1) + (b_2 a'_2 - b'_2 a_2)$ . (Note also that the natural map  $\pi_1(F) \rightarrow H_1(F; Z_2)$  factors through  $G_2$  via  $\psi_2$ .)

**The kernel of  $\psi_2$  is nongeometric** Suppose that there is a simple loop  $\gamma$  representing a nontrivial element  $\omega$  in  $\text{Ker}(\psi_2)$ . Our first observation is that  $\gamma$  can be chosen to be separating. If  $\gamma$  is nonseparating, pick a simple loop  $\gamma'$  meeting  $\gamma$  transversely in exactly one point. Let  $\omega'$  be the element of  $\pi_1(F)$  represented by  $\gamma'$ . Clearly,  $[\omega, \omega']$  is in the kernel of  $\psi_2$  and it is represented by a separating simple loop.

Since  $\gamma$  is now assumed to be separating, it bounds a punctured torus on  $F$ . This follows from the classification of surfaces. Hence  $\omega = [\omega_1, \omega_2]$ , where  $\omega_1$  and  $\omega_2$  are represented by simple loops meeting transversely in one point. From this one computes using (2) that  $\psi_2(\omega) = [\psi_2(\omega_1), \psi_2(\omega_2)] = (0, 0, 0, 0) \times (|\omega_1| \cap |\omega_2|) = (0, 0, 0, 0) \times (1)$ , which is nontrivial in  $G_2$ . This contradicts the assumption that  $\omega \in \text{Ker}(\psi_2)$ .

## 4 Minimality of $G$

The goal of this section is to prove that if  $F$  is of genus 2 and the order of  $G$  is less than 32, then any homomorphism  $\phi: \pi_1(F) \rightarrow G$  has geometric kernel.

Here is a summary of the argument. We first prove that any  $\phi: \pi_1(F) \rightarrow G$  has geometric kernel if  $G$  is a cyclic extension of an abelian group, that is, if  $G$  contains a normal abelian subgroup with cyclic quotient. The approach used to prove this was pointed out by Allan Edmonds. The argument depends on an analysis of the action of the homeomorphism group of  $F$  on the set of representations of  $\pi_1(F)$  to  $G$ . We next note that with the exception of two groups of order 24,  $SL_2(Z_3)$  and  $S_4$ , all groups of order less than 32 are cyclic extensions of abelian groups. This can be proved by a case-by-case analysis using the Sylow theorems. More easily, group tables such as [6] provide the necessary information. The proof is completed using specialized arguments for  $SL_2(Z_3)$  and  $S_4$ .

**Cyclic Extension of Abelian Groups** Fix a group  $G$ . The group of basepoint preserving homeomorphisms of  $F$  acts on the set of representations of  $\pi_1(F)$  to  $G$ , as follows. If  $h$  is a homeomorphism of  $F$ , send a representation  $\phi$  to  $\phi \circ h_*$ . Notice that  $\phi$  has geometric kernel if and only if  $\phi \circ h_*$  has geometric kernel. The following is a result of Nielsen [4]; a proof can be found in [1].

**4.1 Lemma.** *If  $G$  is a cyclic group and  $\phi: \pi(F) \rightarrow G$  is a surjective homomorphism, then there is a homeomorphism  $h$  of  $F$  such that  $\phi \circ h_*(x_1)$  generates  $G$ , and  $\phi \circ h_*(y_1)$ ,  $\phi \circ h_*(x_2)$  and  $\phi \circ h_*(y_2)$  are all trivial.*

**4.2 Theorem.** *If  $G$  contains an abelian normal subgroup  $N$  such that  $G/N$  is cyclic, than any surjective homomorphism  $\phi: \pi_1(F) \rightarrow G$  has a geometric kernel.*

**Proof** Denote the quotient map  $G \rightarrow G/N$  by  $\rho$ . Applying the lemma, we can assume that  $\rho \circ \phi(x_2)$  and  $\rho \circ \phi(y_2)$  are both trivial. Hence  $\phi(x_2)$  and  $\rho \circ \phi(y_2)$  are both in  $N$ . The commutator  $[x_2, y_2]$  is represented by a simple loop and is in the kernel of  $\phi$ , since  $\phi([x_2, y_2])$  is in the commutator subgroup of an abelian group.

## Exceptional Groups

**Case 1** We begin by recalling that  $SL_2(Z_3)$  is isomorphic to the semidirect product of the quaternionic 8-group,  $Q$ , with  $Z_3$ . We will use the standard notation for elements in  $Q$ . The generator of  $Z_3$  will be denoted  $t$ . The action of  $Z_3$  on  $Q$  is given by  $tit^{-1} = j$ ,  $tjt^{-1} = k$  and  $tk t^{-1} = i$ . Note that  $t(-1)t^{-1} = -1$  and that  $-1 \in Q$  is hence central in  $SL_2(Z_3)$ .

Suppose  $\phi: \pi_1(F) \rightarrow SL_2(Z_3)$  is a surjective representation with nongeometric kernel. Applying the lemma to the composition  $\pi_1(F) \rightarrow SL_2(Z_3) \rightarrow SL_2(Z_3)/Q = Z_3$  we can assume that  $\phi(x_1) = tq_1$ ,  $\phi(y_1) = q_2$ ,  $\phi(x_2) = q_3$ , and  $\phi(y_2) = q_4$ , where each  $q_i$  is in  $Q$ .

Since  $[x_2, y_2]$  is represented by a simple loop,  $[q_3, q_4] \neq 1$ . Hence  $[q_3, q_4] = -1 \in Q$ . It follows that  $[tq_1, q_2] = -1$ . Note that  $q_2 \neq \pm 1$ , so  $q_2 = \pm i, \pm j$ , or  $\pm k$ . From the commutator relation,  $tq_1q_2q_1^{-1}t^{-1} = q_2^{-1}$ . For any two quaternions,  $q_1q_2q_1^{-1} = q_2^{\pm 1}$ . Hence,  $tq_2t^{-1} = q_2^{\pm 1}$ . However, this is impossible, given that  $q_2 \neq \pm 1$  and the action of  $t$  on  $Q$ .

**Case 2** The symmetric group  $S_4$  is the semidirect product of  $Z_2 \times Z_2$  with  $S_3$ . As a subgroup, the  $Z_2 \times Z_2$  is given by the set  $\{(1), (12)(34), (13)(24), (14)(23)\}$ . The  $S_3$  is given by the set  $\{(1), (12), (13), (23), (123), (321)\}$ .

Let  $\phi: \pi_1(F) \rightarrow S_4$  be a surjective representation with nongeometric kernel. The main result of [2] applied to the composition  $\pi_1(F) \rightarrow S_4 \rightarrow S_4/(Z_2 \times Z_2) \cong S_3$  shows that by applying a homeomorphism we can arrange that  $\phi$  takes on the values  $\phi(x_1) = (12)n_1$ ,  $\phi(y_1) = n_2$ ,  $\phi(x_2) = (123)n_3$ , and  $\phi(y_2) = n_4$  where each  $n_i$  is in  $Z_2 \times Z_2$ .

Since both  $y_1$  and  $y_2$  are represented by simple loops, neither  $n_2$  nor  $n_4$  are trivial. Also,  $[x_1, y_1]$  is represented by a simple loop, so  $[(12)n_1, n_2] \neq 1$ . It follows that  $n_2 \neq (12)(34)$ . There are two other possibilities for  $n_2$ .

Suppose that  $n_2 = (13)(24)$ . There are three possible values of  $n_4$  to be considered. Because  $y_1y_2$  is realized by a simple loop,  $n_4 \neq (13)(24)$ . It is easily seen that  $y_1x_2y_2^{-1}x_2^{-1}$  is realized by a simple loop. Hence  $n_4 \neq (12)(34)$ . Finally  $n_4 \neq (14)(23)$ , because  $y_2x_1y_1^{-1}x_1^{-1}$  can also be represented by a simple loop.

We proceed similarly if  $n_2 = (14)(23)$ . Clearly  $n_4 \neq (14)(23)$ . Because  $y_2x_1y_1^{-1}x_1^{-1}$  is realized by a simple loop,  $n_4 \neq (13)(24)$ . Finally, it is again easily seen that  $y_1x_2^{-1}y_2^{-1}x_2$  is realized by a simple loop. This implies that  $n_4 \neq (12)(34)$ . All possibilities have now been eliminated.

## 5 Generalizations

The group constructed in Section 3,  $G_2$ , can be generalized to a group  $G_k$  such that for the genus  $k$  surface  $F_k$  there is a homomorphism  $\phi_k: \pi_1(F_k) \rightarrow G_k$  with nongeometric kernel. The arguments are similar to those of Section 3 and are only outlined here.

Define  $G_k$  by defining a product on the set  $(Z_k)^{2k} \times Z_k$  as follows.

$$(a_1, b_1, a_2, b_2, \dots, b_k, \epsilon)(a'_1, b'_1, a'_2, b'_2, \dots, b'_k, \epsilon') = (a_1 + a'_1, b_1 + b'_1, a_2 + a'_2, b_2 + b'_2, \dots, b_k + b'_k, \epsilon + \epsilon' + \sum b_i a'_i)$$

Sums and products within the parenthesis are in  $Z_k$ . That this defines a group is a straightforward calculation.

There is a natural representation  $\phi_k: \pi_1(F_k) \rightarrow G_k$  as before. In this case the essential observation is

$$[\phi_k(\omega_1), \phi_k(\omega_2)] = (0, 0, \dots, 0) \times (|\omega_1| \cap |\omega_2|), \quad (1)$$

where  $(|\omega_1| \cap |\omega_2|)$  is the  $Z_k$  intersection number of the classes in  $H_1(F_k; Z_k)$  represented by  $\omega_1$  and  $\omega_2$ .

If  $\phi_k$  had geometric kernel, there would be a separating simple loop representing an element in the kernel. Using the classification of surfaces, that element would be of the form

$$[\omega_1, \omega'_1][\omega_2, \omega'_2] \cdots [\omega_m, \omega'_m]$$

with  $m < k$  and  $(|\omega_i| \cap |\omega'_i|) = 1$  for all  $i$ . A contradiction follows as in Section 3.

**Remark** The order of the group  $G$  just constructed is  $g^{2g+1}$ . This number should be contrasted to the order found in Section 2,  $2^{(g-1)2^{2g+1}+2+2g}$ . The first is obviously much smaller than the second. The results of this paper, along with our difficulties in trying to find smaller examples, leads us to conjecture that  $g^{2g+1}$  represents the least possible order.

### References

- [1] Edmonds, A. *Surface Symmetry I*, Michigan J. Math 29 (1982) 171–183.
- [2] Edmonds, A. *Surface Symmetry II*, Michigan J. Math 30 (1983) 143–154.
- [3] Hall, M. and Senior, J.K. *The groups of order  $2^n$  ( $n \leq 6$ )* Macmillan, New York (1964).
- [4] Nielson, J. *Die Struktur periodischer Transformationen von Flächen*, Dansk Vid. Selsk., Mat.-Fys. Medd. 15 (1937), 1–77.
- [5] Skora, R. *Dissertation*, Department of Mathematics, University of Texas, Austin, Texas, 1984.
- [6] Thomas, A.D. and Wood, G.V. *Group Tables*, Shiva Publishing Limited, Kent, Great Britian, 1980.

Department of Mathematics  
Indiana University  
Bloomington, IN 47405  
livingst@indiana.edu